arXiv:hep-ex/0008061v1 24 Aug 2000

DPF '96: The Minneapolis Meeting (11-15 August 1996) pages 868–870.

$D-\pi$ Production Correlations in 500 GeV $\pi^- - N$ Interactions

K. GOUNDER and L. CREMALDI
for the Fermilab E791 Collaboration
Dept. of Physics and Astronomy, University of Mississippi, Oxford, MS 38677

We study the charge correlations between charm mesons produced in 500 GeV $\pi^- - N$ interactions and the charged pions produced closest to them in phase space. With 110,000 fully reconstructed D mesons from experiment E791 at Fermilab, the correlations are studied as functions of the $D\pi - D$ mass difference and of Feynman x. We observe significant correlations which appear to originate from a combination of sources including fragmentation dynamics, resonant decays, and charge of the beam.

In the production of a D meson the charge of pions formed nearby in phase space may be highly correlated with the charge of the D meson. For example, this *associated pion* can be produced in the fragmentation dynamics[1] of a c quark combining with a $\overline{d}$ from a $d-\overline{d}$ vacuum pair, forming a D^+, and the remaining d forming an associated pion nearby in phase space. Thus, $D^+(D^{*+})\pi^-$, $D^-(D^{*-})\pi^+$, and $D^0\pi^+$ $\overline{D^0}\pi^-$ would be favored over combinations with opposite pion charges. Resonant decays[2] and the charge of the beam particle are also sources of charge correlations. Using a π^- beam can lead to the association of both charm mesons and anticharm mesons with negative pions, especially in the forward (beam) direction. By comparing the charge correlations of different species of charm mesons and antimesons with associated pions, and by studying them as functions of Feynman x (x_F), one can hope to disentangle some of these processes.

We use well identified $D^0 \to K^-\pi^+$, $D^+ \to K^-\pi^+\pi^+$, and $D^{*+} \to D^0\pi^+$ signals (and their charge conjugate decays) from experiment E791 at Fermilab for this study. A silicon vertex detector and two Čerenkov counters are used to locate decay vertices and identify pions and kaons for this analysis. Details of the spectrometer and event selection are found elsewhere.[3] The final D meson sample sizes used for this analysis are 22587 ± 210, 24237 ± 216, 24569 ± 204, 29649 ± 238, 4997 ± 84, and 6048 ± 93 events, for D^0, $\overline{D^0}$, D^+, D^-, D^{*+}, and D^{*-}, respectively.

For each D found in an event, all identified pion tracks originating from the primary vertex are combined with the D. Among these combinations, the pion that forms the smallest invariant mass difference (Δm_{min}) with the D is selected as the candidate associated pion.

Table 1: The x_F- and Δm-integrated correlation parameters α defined in Eq. (1) for the background-subtracted signals prior to correction, for the corresponding backgrounds, and for the signals after correction using the matrix correction technique.

Charm	Signal α	Background α	Corrected Signal α
D^0	0.13 ± 0.01	-0.04 ± 0.01	$0.12 \pm 0.03 \pm 0.04$
$\overline{D^0}$	0.18 ± 0.01	0.04 ± 0.01	$0.42 \pm 0.02 \pm 0.03$
D^+	0.18 ± 0.01	0.10 ± 0.01	$0.45 \pm 0.03 \pm 0.03$
D^-	0.08 ± 0.01	0.02 ± 0.01	$0.03 \pm 0.03 \pm 0.04$
D^{*+}	0.15 ± 0.02	0.08 ± 0.03	$0.33 \pm 0.05 \pm 0.03$
D^{*-}	0.08 ± 0.02	0.02 ± 0.03	$0.15 \pm 0.05 \pm 0.04$

We define the correlation parameter α as

$$\alpha(D) \equiv \frac{\sum N_i(D\pi^q) - \sum N_i(D\pi^{-q})}{\sum N_i(D\pi^q) + \sum N_i(D\pi^{-q})} \tag{1}$$

where $q = +, -, -, +, -, +$ for $D = D^0$, $\overline{D^0}$, D^+, D^-, D^{*+}, and D^{*-}, respectively, and $\sum N_i(D\pi^q)$ denotes the number of charm mesons for which the selected pion has the charge q.

The correlation parameters for background-subtracted signals and background regions are listed in Table 1. The signal correlations differ significantly from the background correlations. These raw correlations must be corrected for the occurrence of falsely reconstructed tracks (ghosts), momentum smearing in bins of Δm_{min}, detector acceptance, and dilution of the correlation by wrong track selection. We use a Monte Carlo simulation of the experiment and the LUND event generator (PYTHIA 5.7/JETSET 7.3)[4] to model these effects and extract the final results (Table 1), within a matrix formalism explained elsewhere.[5] A fundamentally different model of hadron production could change the corrections; an effect not accounted for in Table 1.

To investigate beam-related effects in more detail, we study the x_F dependence of the D^+ and D^{*+} correlations. In Fig. 1, we plot α as a function of x_F, for both particle and antiparticle for D^+ and D^{*+}. We observe that $\alpha(D^+)$ rises slightly with x_F but $\alpha(D^-)$ falls sharply to negative values for $x_F > 0.2$. In both cases, the D's are more likely to be associated with π^-'s at high x_F where beam effects seem to be important. There appears to be less dependence of α on x_F for the $D^{*\pm}$.

Detailed Monte Carlo studies suggest minimal or no beam-related effects when the combined particle and antiparticle correlations are computed. We calculate the combined correlation parameters to be $\alpha(D^0, \overline{D^0}) = 0.29 \pm 0.02 \pm 0.03$, $\alpha(D^+, D^-) = 0.21 \pm 0.02 \pm 0.03$, and $\alpha(D^{*+}, D^{*-}) = 0.23 \pm 0.04 \pm 0.03$. These results indicate that fragmentation dynamics and resonant decays produce substantial correlations between D mesons and their associated pions. All three combined correlation levels are approximately equal.

In summary, we observe significant production correlations between D mesons and their associated pions. Beam-related and other fragmentation effects play an important role. Further studies are needed to unravel the contributions.

This work was supported in part by U.S. DOE DE–FG05–91ER40622.

1. M. Gronau and J. L. Rosner, Phys. Rev. D **49**, 254 (1994).
2. M. Gronau, A. Nippe, and J. L. Rosner, Phys. Rev. D **47**, 1988 (1992).
3. E791 Collaboration, Aitala et al., Phys. Lett. B **371**, 157 (1996).
4. T. Sjöstrand, Comput. Phys. Commun. **82**, 74 (1994).
5. E791 Collaboration, Aitala et al., Fermilab-Pub-96/206-E and Phys. Lett. B **403**, 185 (1997).



Figure 1: The corrected correlation parameter α as a function of x_F for (a)D^+ and (b)D^{*+}. The parameter α is defined in Eq. (1) in the text. The error bars correspond to the statistical and systematic uncertainties added in quadrature.